

Lecture 3

Conditional Probability & Independence

Manju M. Johny

STAT 330 - Iowa State University

Contingency Table

Contingency Table

Definition

A *contingency table* gives the distribution of 2 variables.

Example 1: Suppose in a small college of 1000 students, 650 students own iPhones, 400 students own MacBooks, and 300 students own both.

Define events: I = “owns iPhone”, and M = “owns MacBook”.

Phone \ Computer	M	\bar{M}	Total
	I	300	?
\bar{I}	?	?	?
Total	400	?	1000

Contingency Table

Phone \ Computer	M	\bar{M}	Total
	I	300	350
\bar{I}	100	250	350
Total	400	600	1000

Marginal Probability

Marginal Probability

Definition

The *marginal probability* is the probability of a variable. It can be obtained from the *margins* of contingency table.

Phone \ Computer	M	\bar{M}	Total
	I	300	350
\bar{I}	100	250	350
Total	400	600	1000

What is the probability of owning a Mac? (ie marginal probability of owning a Mac)

$$P(M) = \frac{400}{1000} = 0.40$$

Conditional Probability

Conditional Probability

Does knowing someone owns an Iphone change the probability they own a Mac?

Informally, conditional probability is updating the probability of an event given information about another event.

If we *know* that someone owns an Iphone, then we can narrow our sample space to just the “owns Iphone” case (highlighted blue row) and ignore the rest!

Phone \ Computer	Computer		Total
	M	\bar{M}	
I	300	350	650
\bar{I}	100	250	350
Total	400	600	1000

Conditional Probability Cont.

What is the probability of owning a Mac *given* they own an Iphone?

		Computer		Total
		M	\bar{M}	
Phone	I	300	350	650
	\bar{I}	100	250	350
Total		400	600	1000

$$P(M|I) = \frac{300}{650} = 0.46$$

Conditional Probability Cont.

Definition

The *conditional probability* of event A given event B is

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

provided $P(B) \neq 0$.

It can be obtained from the *rows/columns* of contingency table.

Back to [Example 1](#) ...

What is the probability of owning a Mac *given* they own an Iphone?

$$P(M|I) = \frac{P(I \cap M)}{P(I)} = \frac{0.3}{0.65} = 0.46$$

Consequences of Conditional Probability

The definition of conditional probability gives useful results:

1.

$$P(A|B) = \frac{P(A \cap B)}{P(B)} \rightarrow P(A \cap B) = P(B)P(A|B)$$

2.

$$P(B|A) = \frac{P(A \cap B)}{P(A)} \rightarrow P(A \cap B) = P(A)P(B|A)$$

This gives us two additional ways to calculate probability of intersections. Putting it together ...

$$P(A \cap B) = P(A|B)P(B) = P(B|A)P(A)$$

Probability Calculations

Probability Calculations

A contingency table can also be written with probabilities instead of counts. This is called a *probability table*.

Inner cells give “joint probabilities” → probability of intersections

- $P(A \cap B), P(\bar{A} \cap B)$, etc

Margins give “marginal probabilities” → probability of variables

- $P(A), P(B), P(\bar{A})$, etc

Phone \ Computer	M	\bar{M}	Total
	I	0.30	0.35
\bar{I}	0.10	0.25	0.35
Total	0.40	0.60	1

Probability Calculations Cont.

Phone \ Computer	M	\bar{M}	Total
	I	0.30	0.35
\bar{I}	0.10	0.25	0.35
Total	0.40	0.60	1

$$P(\bar{I}) =$$

$$P(M) =$$

$$P(\bar{I} \cap M) =$$

$$P(M|\bar{I}) =$$

$$P(\bar{I}|M) =$$

Independence

Independence of Events

In Example 1, knowing an event occurred changed the probability of another event occurring.

However, sometimes knowing an event occurs *doesn't change* the probability of the other event.

In this case, we say the events are *independent*.

Definition

Events A and B are *independent* if ...

1. $P(A \cap B) = P(A)P(B)$

or equivalently

2. $P(A|B) = P(A)$ if $P(B) \neq 0$

Independence of Events Cont.

Example 2: Check if events are independent

Is owning an Iphone and owning MacBook independent?

Recall that $P(I) = 0.65$, $P(M) = 0.4$, $P(I \cap M) = 0.35$

Independence of Events Cont.

Example 3: Using independence to simplify calculations

If A, B independent $\rightarrow P(A \cap B) = P(B)P(A|B) = P(B)P(A)$

Roll a die 4 times. Assuming that rolls are independent, what is the probability of obtaining at least one '6'?

$$P(\text{at least 1 '6'}) = 1 - P(\text{No '6's})$$

$$= 1 - P(\text{no '6' on roll 1} \cap \text{no '6' on roll 2} \cap \dots \cap \text{no '6' on roll 4})$$


=

Independent vs. Disjoint

Independent \neq Disjoint!!!


Completely different concepts!

Independent:



$$P(A \cap B) = P(A)P(B)$$

Disjoint:




$$P(A \cap B) = P(\emptyset) = 0$$

System Reliability

Application: System Reliability

Parallel: A parallel system consists of k components (c_1, \dots, c_k) arranged such that the system works if and only if at least one of the k components functions properly.



Series: A series system consists of k components (c_1, \dots, c_k) arranged such that the system works if and only if ALL components function properly.



Reliability: Reliability of a system is the probability that the system works.

Reliability of Parallel System

Example 4:

Let c_1, \dots, c_k denote the k components in a *parallel* system. Assume the k components operate independently, and $P(c_j \text{ works}) = p_j$. What is the reliability of the system?

$$\begin{aligned} P(\text{system works}) &= P(\text{at least one component works}) \\ &= 1 - P(\text{all components fail}) \\ &= 1 - P(c_1 \text{ fails} \cap c_2 \text{ fails} \cap \dots \cap c_k \text{ fails}) \\ &= 1 - \prod_{j=1}^k P(c_j \text{ fails}) \\ &= 1 - \prod_{j=1}^k (1 - p_j) \end{aligned}$$

Reliability of Series System

Example 5:

Let c_1, \dots, c_k denote the k components in a *series* system. Assume the k components operate independently, and $P(c_j \text{ works}) = p_j$. What is the reliability of the system?

$$\begin{aligned}P(\text{system works}) &= P(\text{all components work}) \\&= P(c_1 \text{ works} \cap c_2 \text{ works} \cap \dots \cap c_k \text{ works}) \\&= \prod_{j=1}^k P(c_j \text{ works}) \\&= \prod_{j=1}^k p_j\end{aligned}$$

Reliability Example

Example 6: Suppose a base is guarded by 3 radars (R_1, R_2, R_3), and the radars are independent of each other. The detection probability are ...

$$P(R_1 \text{ detects}) = 0.95$$

$$P(R_2 \text{ detects}) = 0.98$$

$$P(R_3 \text{ detects}) = 0.99$$

Does a system in *parallel* or *series* have higher reliability for this scenario?

Reliability Example

Reliability Example